

Mucinous Adenocarcinoma of the Urinary Bladder after Long-Term Duodeno-Renal and Colovesical Fistula - Case Report

J. Petrovic¹, G. Barisic¹, Z. Krivokapic¹, B. Krivokapic²

¹Department for Colorectal Diseases, Clinic for Digestive Surgery, 1st Surgical Clinic, Clinical Center of Serbia, Belgrade, Serbia School of Medicine, University of Belgrade, Serbia

²Orthopedic Clinic Banjica, School of Medicine University of Belgrade, Serbia

Abstract

Primary adenocarcinoma of the urinary bladder is a rare neoplasm. It accounts for 1-2% of all bladder carcinomas and sometimes may be found in the bladder diverticula. Fistula between duodenum and renal pelvis is another rarity while colovesical fistula is not so uncommon. We present a case of a 40 years old man who had surgery for colovesical and duodenorenal fistula and subsequently developed adeno-carcinoma of the urinary bladder.

Key words: duodeno-pyelic fistula, sigmoido-urethral fistula, adenocarcinoma, urinary bladder

Corresponding author: Zoran V. Krivokapic, MD, FRCS

First Surgical Clinic

Koste Todorovica 6, 11000 Belgrade, Serbia

Tel/fax: +381 11 3622 811

E-mail: scpy@beotel.rs